

# Drug Info

## ACETAMIN. + BUTALBIT. + CAFF. (FIORICET)

### BEFORE USING THIS MEDICINE:

- \* Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- \* Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.
- \* Inform your doctor if you have any of these conditions : LIVER PROBLEMS; If you are PREGNANT.

### COMMON USES FOR THIS MEDICINE:

- 1- For the treatment of TENSION HEADACHE
- 2- For the treatment of PAIN, GENERALIZED
- 3- For the treatment of PYREXIA OF UNKNOWN ORIGIN

### HOW TO USE THIS MEDICINE:

- Use this medicine exactly as directed on the label, unless instructed differently by your doctor
- \* This medicine can be taken with or without food, REGARDLESS of meal times. If stomach upset occurs, take with food.

### WHAT TO DO IF YOU MISS A DOSE:

Take when remembered unless time for next dose. No double doses.

### HOW TO STORE THIS MEDICINE:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

### CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Yellowing of eyes or skin; Dizziness; Irregular heartbeat; Swelling usually of face or lips; Skin rash; Skin inflammation; Severe skin reactions.

### OTHER SIDE EFFECTS:

Drowsiness; Lightheadedness; Nausea; Slowness of heartbeat; Sleepiness; Stomach discomfort or pain; Low blood pressure; Anemia; Abnormal blood cell count; Hallucinations; Confusion; Constipation; Depression; Headache; Nightmares; Nervousness; Vomiting.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

### PRECAUTIONS AND WARNINGS:

- \* There are no adequate or well-controlled safety studies in pregnant women. NOTIFY YOUR DOCTOR IF PREGNANT or intend to become pregnant.
- \* This medicine may cause drowsiness, dizziness, or blurred vision. DO NOT drive or operate machines until you know how you react to this medicine.
- \* AVOID concurrent use of alcohol or other CNS depressants as it may intensify the drowsiness, dizziness, or lightheadedness effect of this medicine.
- \* Strictly limit alcohol consumption. LIVER PROBLEMS may be more likely to occur if you drink alcoholic beverages regularly while taking this medicine.
- \* Inform MD of other medications you are taking concurrently including over-the-counter medicines and herbal or dietary supplements.
- \* Prolonged use of this medication may cause physical and psychological dependence, do not exceed recommended dose.