

Drug Info

AMOXICILLIN + POT CLAVUL (AUGMENTIN ES)

BEFORE USING THIS MEDICINE:

- * Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- * Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.
- * Inform your doctor if you have any of these conditions : LIVER PROBLEMS; KIDNEY DISEASE.

COMMON USE FOR THIS MEDICINE:

1- For the treatment of recurrent episodes of OTITIS MEDIA (SUPPURATIVE AND UNSPECIFIED)

HOW TO USE THIS MEDICINE:

Use this medicine exactly as directed on the label, unless instructed differently by your doctor

- * Use this medicine for the full prescribed treatment period.
- * SHAKE suspension well before using.
- * Take this medicine WITH food, a meal, or a snack. Do not take on an empty stomach.

WHAT TO DO IF YOU MISS A DOSE:

Take when remembered unless time for next dose. No double doses.

HOW TO STORE THIS MEDICINE:

Refrigerate in original container away from children. Do not freeze. Discard unneeded medicine.

STOP TAKING THIS MEDICINE AND GET EMERGENCY HELP IMMEDIATELY IF ANY OF THE FOLLOWING OCCURS (SEE SYMPTOMS BELOW):

Shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching.

CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Skin rash; Unusual bleeding or bruising; Skin inflammation; Diarrhea; Fungal infection of mouth.

OTHER SIDE EFFECTS:

Headache; Nausea; Black tongue; Vomiting.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

PRECAUTIONS AND WARNINGS:

- * STOP TAKING MEDICINE & GET EMERGENCY HELP IMMEDIATELY IF A HYPERSENSITIVITY OR ANAPHYLACTIC REACTION occurs: (see symptoms below):-
 - * shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching.
 - * There are no adequate or well-controlled safety studies in pregnant women. NOTIFY YOUR DOCTOR IF PREGNANT or intend to become pregnant.
- * TELL YOUR DOCTOR if you have ever had any unusual or ALLERGIC reaction to any of the CEPHALOSPORINS, PENICILLINS, or penicillin-like medicines.
- * To prevent drug resistance, this medication should be used to treat BACTERIAL INFECTIONS ONLY. It is not effective against viral infections.
- * This medicine may cause false test results with some urine tests. If you are DIABETIC, check with your doctor before changing diabetes regimen.
- * If DIARRHEA develops/persists and is severe or lasts more than 2 or 3 days, call your doctor.

NUTRITIONAL SUPPLEMENTS FOR DEPLETION:

This medication may cause some nutrients to be depleted from your body. The following supplements may be recommended:

BIOTIN FORTE; INOSITOL ORAL; LACTOBACILLUS ORAL (LACTINEX); VITAMIN B1 ORAL (THIAMINE); VITAMIN B2 ORAL (RIBOFLAVIN); VITAMIN B3 ORAL (NICOTINAMIDE); VITAMIN B6 ORAL (PYRIDOXINE); VITAMIN B12 ORAL; PHYTONADIONE-VIT K1 ORAL (MEPHYTON); VITAMIN K.