

Palmetto State Pharmaceuticals

Patient Drug Education

AMOXICILLIN 500MG CAPSULE

USES:

Amoxicillin is a penicillin-type antibiotic used to treat a wide variety of bacterial infections. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections. It will not work for viral infections (e.g., common cold, flu). Unnecessary use or overuse of any antibiotic can lead to its decreased effectiveness.

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Take this medication by mouth with or without food, usually every 8 or 12 hours, or as directed by your doctor. The dosage is based on your medical condition and response to therapy. Drink plenty of fluids while using this medication unless your doctor tells you otherwise. Antibiotics work best when the amount of medicine in your body is kept at a constant level. Therefore, take this drug at evenly spaced intervals. Continue to take this medication until the full-prescribed amount is finished even if symptoms disappear after a few days. Stopping the medication too early may allow bacteria to continue to grow, which may result in a relapse of the infection. Inform your doctor if your condition persists or worsens.

SIDE EFFECTS:

Nausea, vomiting or diarrhea may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: dark urine, persistent nausea or vomiting, stomach/abdominal pain, yellowing eyes or skin, easy bruising or bleeding, persistent sore throat or fever. This medication may rarely cause a severe intestinal condition (pseudomembranous colitis) due to a resistant bacteria. This condition may occur weeks after treatment has stopped. Do not use anti-diarrhea products or narcotic pain medications if you have the following symptoms because these products may make them worse. Tell your doctor immediately if you develop: persistent diarrhea, abdominal or stomach pain/cramping, or blood/mucus in your stool. Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection (oral or vaginal fungal infection). Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge or other new symptoms. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. Amoxicillin can commonly cause a mild rash that is usually not serious. However, you may not be able to tell it apart from a rare rash that could be a sign of a severe allergic reaction. Therefore, seek immediate medical attention if you develop any rash. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking amoxicillin, tell your doctor or pharmacist if you are allergic to it; or to penicillin or cephalosporin antibiotics; or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, a certain type of viral infection (infectious mononucleosis). Kidney function declines as you grow older. This medication is removed by the kidneys. Therefore, elderly people may be more sensitive to this drug. This medication should be used only when clearly needed during pregnancy. Discuss the risks and benefits with your doctor. Amoxicillin passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: allopurinol, live bacterial vaccines, methotrexate, tetracyclines. Before taking amoxicillin, tell your doctor or pharmacist if you are also taking probenecid. Probenecid slows down the removal of amoxicillin from your body, resulting in higher levels of this antibiotic in your bloodstream. For certain types of difficult-to-treat infections, your doctor may prescribe these 2 medications together in order to achieve this effect. Consult your doctor or pharmacist for more details. This medication may decrease the effectiveness of combination-type birth control pills. This can result in pregnancy. You may need to use an additional form of reliable birth control while using this medication. Consult your doctor or pharmacist for details. Amoxicillin may cause false positive results with certain diabetic urine testing products (cupric sulfate-type). This drug may also affect the results of certain lab tests. Make sure laboratory personnel and your doctors know you use this drug. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES:

Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless told to do so by your doctor. A different medication may be necessary in those cases. With prolonged treatment, laboratory and/or medical tests (e.g., kidney and liver function, complete blood counts) should be performed periodically to monitor your progress or check for side effects. Consult the doctor for more details.

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

Palmetto State Pharmaceuticals

STORAGE:

Store at room temperature according to the product labeling, away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets.