

Palmetto State Pharmaceuticals

Patient Drug Education

AUGMENTIN 500-125 TABLET

USES:

Amoxicillin/clavulanic acid is a penicillin-type antibiotic used to treat a wide variety of bacterial infections. It works by stopping the growth of bacteria. This antibiotic only treats bacterial infections. It will not work for viral infections (e.g., common cold, flu). Unnecessary use or overuse of any antibiotic can lead to its decreased effectiveness.

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Take this medication by mouth with a meal or snack, usually every 8 or 12 hours, or as directed by your doctor. Antibiotics work best when the amount of medicine in your body is kept at a constant level. Therefore, take this drug at evenly spaced intervals. Continue to take this medication until the full prescribed amount is finished even if symptoms disappear after a few days. Stopping the medication too early may allow bacteria to continue to grow, which may result in a relapse of the infection. Inform your doctor if your condition persists or worsens.

SIDE EFFECTS:

Diarrhea, nausea, or vomiting may occur during the first few days as your body adjusts to the medication. Take with food to minimize stomach upset. If any of these effects persist or worsen, contact the doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: easy bruising or bleeding, persistent sore throat or fever, dark urine, persistent nausea or vomiting, severe stomach/abdominal pain, yellowing eyes or skin. This medication may rarely cause a severe intestinal condition (pseudomembranous colitis) due to a resistant bacteria. This condition may occur weeks after treatment has stopped. Do not use anti-diarrhea products or narcotic pain medications if you have the following symptoms because these products may make them worse. Tell your doctor immediately if you develop: persistent diarrhea, severe abdominal or stomach pain/cramping, or blood/mucus in your stool. Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection (oral or vaginal fungal infection). Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge or other new symptoms. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. The amoxicillin in this combination medication can commonly cause a mild rash that is usually not serious. However, you may not be able to tell it apart from a rare rash that could be a sign of a severe allergic reaction. Therefore, seek immediate medical attention if you develop any rash. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking amoxicillin/clavulanic acid, tell your doctor or pharmacist if you are allergic to it; or to penicillin or cephalosporin antibiotics; or if you have any other allergies. This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: a history of liver problems (e.g., cholestatic jaundice) associated with prior use of amoxicillin/clavulanic acid. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease, a certain type of viral infection (infectious mononucleosis). Kidney function declines as you grow older. This medication is removed by the kidneys. Therefore, elderly people may be more sensitive to this drug. This medication should be used only when clearly needed during pregnancy. Caution is advised if this medication is given before delivery for a certain condition (premature rupture of fetal membranes) due to a possible increased risk of harm to the newborn. Discuss the risks and benefits with your doctor. This medication passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Before using this medication, tell the doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: allopurinol, live bacterial vaccines, methotrexate, tetracyclines. Before taking amoxicillin/clavulanic acid, tell your doctor or pharmacist if you are also taking probenecid. Probenecid slows down the removal of amoxicillin/clavulanic acid from your body, resulting in higher levels of this antibiotic in your bloodstream. For certain types of difficult-to-treat infections, your doctor may prescribe these 2 medications together in order to achieve this effect. Consult your doctor or pharmacist for more details. This medication may decrease the effectiveness of combination-type birth control pills. This can result in pregnancy. You may need to use an additional form of reliable birth control while using this medication. Consult your doctor or pharmacist for details. This antibiotic may cause false positive results with certain diabetic urine testing products (cupric sulfate-type). This drug may also affect the results of certain lab tests. Make sure laboratory personnel and your doctors know you use this drug. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES:

Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection. A different medication may be necessary in those cases. With prolonged treatment, laboratory and/or medical tests (e.g., kidney and liver function, complete blood counts) should be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

Palmetto State Pharmaceuticals

MISSED DOSE:

If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store at or below room temperature (77 degrees F or 25 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets.