

# Palmetto State Pharmaceuticals

---

## *Patient Drug Education*

---

---

### **AZITHROMYCIN 200 MG/5 ML SUSP**

---

#### **USES:**

Azithromycin is an antibiotic (macrolide-type) used to treat a wide variety of bacterial infections. This antibiotic only treats bacterial infections. It will not work for viral infections (e.g., common cold, flu). Unnecessary use or overuse of any antibiotic can lead to its decreased effectiveness.

#### **HOW TO TAKE THIS MEDICATION:**

**HOW TO USE:** Shake the suspension well before using. Take this medication by mouth with or without food, usually once a day, or as directed by your doctor. You may take this medication with food if stomach upset occurs. The dosage is based on your medical condition and response to therapy. If you are giving this medication to a child, the dosage is also based on his/her weight. Antibiotics work best when the amount of medicine in your body is kept at a constant level. Therefore, take this drug at the same time each day if taken daily. Continue to take this medication until the full prescribed amount is finished even if symptoms disappear after a few days. Stopping the medication too early may allow bacteria to continue to grow, which may result in a relapse of the infection. Antacids may decrease the absorption of azithromycin. If you take an antacid, wait at least 2 hours after taking this medication. Inform your doctor if your condition persists or worsens.

#### **SIDE EFFECTS:**

Stomach upset, diarrhea/loose stools, nausea, vomiting, or stomach/abdominal pain may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: hearing loss. Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: dark urine, persistent nausea/vomiting, severe stomach/abdominal pain, yellowing of the eyes or skin. This medication may rarely cause a severe intestinal condition (pseudomembranous colitis) due to a resistant bacteria. This condition may occur weeks after treatment has stopped. Do not use anti-diarrhea products or narcotic pain medications if you have the following symptoms because these products may make them worse. Tell your doctor immediately if you develop: persistent diarrhea, abdominal or stomach pain/cramping, or blood/mucus in your stool. Use of this medication for prolonged or repeated periods may result in oral thrush or a new vaginal yeast infection (oral or vaginal fungal infection). Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge or other new symptoms. A serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction include: rash, itching, swelling, dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

#### **PRECAUTIONS:**

Before taking azithromycin, tell your doctor or pharmacist if you are allergic to it; or to other macrolide antibiotics such as erythromycin, clarithromycin; or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease. This medication should be used only when clearly needed during pregnancy. Discuss the risks and benefits with your doctor. This drug passes into breast milk. Consult your doctor before breast-feeding.

#### **DRUG INTERACTIONS:**

See also the How to Use section. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: aluminum- and magnesium-containing antacids, digoxin, live bacterial vaccines, lovastatin, nelfinavir, warfarin. This medication may decrease the effectiveness of combination-type birth control pills. This can result in pregnancy. You may need to use an additional form of reliable birth control while using this medication. Consult your doctor or pharmacist for details. Other drugs besides azithromycin which may affect the heart rhythm (QTc prolongation in the EKG) include dofetilide, pimozide, procainamide, quinidine, sotalol, propafenone, and sparfloxacin among others. QTc prolongation can infrequently result in serious, rarely fatal, irregular heartbeats. Consult your doctor or pharmacist for details. Ask for instructions about whether you need to stop any other QTc-prolonging drugs you may be using in order to minimize the risk of this effect. Do not start or stop any medicine without doctor or pharmacist approval.

#### **NOTES:**

Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless told to do so by your doctor. A different medication may be necessary in those cases.

#### **MISSED DOSE:**

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

#### **STORAGE:**

# Palmetto State Pharmaceuticals

Store between 41-86 degrees F (5-30 degrees C) away from light and moisture. Do not store in the bathroom. Discard after 10 days. Keep all medicines away from children and pets.