

# Palmetto State Pharmaceuticals

---

## *Patient Drug Education*

---

---

### **BETAMETHASONE DP 0.05% CRM**

---

#### **USES:**

This medication is used to treat a variety of skin conditions (e.g., eczema, dermatitis, allergies, rash). Betamethasone reduces the swelling, itching, and redness that can occur in these types of conditions. This medication is a strong corticosteroid.

#### **HOW TO TAKE THIS MEDICATION:**

**HOW TO USE:** Use this medication on the skin only. However, do not use it on the face, groin, or underarms or for diaper rash unless directed to do so by your doctor. Wash and dry your hands. Before applying the medication, clean and dry the affected area. Apply a thin film of the medication to the affected area and gently rub in, usually 1-2 times daily or as directed by your doctor. Do not cover, bandage, or wrap the area unless directed to do so by your doctor. If your doctor directs you to use this medication in the diaper area on an infant, do not use tight-fitting diapers or plastic pants. After applying the medication, wash your hands unless you are using this medication to treat the hands. When applying this medication near the eyes, avoid getting it in the eyes as this may worsen or cause glaucoma. Also, avoid getting this medication in the nose or mouth. If you get the medication in these areas, rinse with plenty of water. Use this medication only for the condition for which it was prescribed. Do not use it for longer than prescribed. Inform your doctor if your condition persists or worsens after 2 weeks.

#### **SIDE EFFECTS:**

Burning, itching, irritation, or dryness at the application site may occur when this medication is first applied to the skin. This should disappear in a few days as your body adjusts to the medication. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because he or she has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor promptly if any of these unlikely but serious side effects occur: stretch marks, skin thinning/discoloration, acne, extreme/unwanted hair growth, "hair bumps" (folliculitis). Skin infections can become worse when this medication is used. Notify your doctor if redness, swelling, or irritation does not improve. A very serious allergic reaction to this drug is rare. However, seek immediate medical attention if you notice any symptoms of a serious allergic reaction, including: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

#### **PRECAUTIONS:**

Before using betamethasone, tell your doctor or pharmacist if you are allergic to it; or to other corticosteroids (e.g., hydrocortisone, prednisone); or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: poor blood circulation, immune system problems. Do not use if there is an infection or sore present in the area to be treated. Though very unlikely, it is possible this medication will be absorbed into your bloodstream. This may have undesirable consequences that may require additional corticosteroid treatment. This is especially true for children, those who have used this drug for an extended period of time, and those who have serious medical problems such as serious infections, injuries, or surgeries. This precaution applies for up to one year after you stop using this drug. Tell your doctor immediately if any of the following side effects occur: vision problems, persistent headache, increased thirst/urination, unusual weight loss, unusual weakness, dizziness. Consult your doctor or pharmacist for more details and inform them that you use or have used this medication. Caution is advised when using this drug in children because they may be more sensitive to the effects of too much steroid hormone. Though it is unlikely to occur with corticosteroids applied to the skin, this medication may affect growth in infants and children if used for prolonged periods. Monitor your child's height and rate of growth periodically. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is not known whether this drug passes into breast milk when applied to the skin. Similar medications pass into breast milk when taken by mouth. Consult your doctor before breast-feeding.

#### **DRUG INTERACTIONS:**

Your doctor or pharmacist may already be aware of any possible drug interactions and may be monitoring you for them. Do not start, stop, or change the dosage of any medicine before checking with your doctor or pharmacist first. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: corticosteroids taken by mouth (e.g., prednisone), drugs that lower the immune system (e.g., cyclosporine).

#### **NOTES:**

Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for other skin problems unless told to do so by your doctor. A different medication may be necessary in those cases. Laboratory and/or medical tests (e.g., adrenal gland function tests, morning cortisol blood test) may be performed periodically to monitor your progress or check for side effects, especially if you use this drug for an extended time, apply it over large areas of the body, or bandage/wrap the treated areas. Consult your doctor for more details. Inform all your doctors you use or have used this medication.

#### **MISSED DOSE:**

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule.

# Palmetto State Pharmaceuticals

Do not double the dose to catch up.

## **STORAGE:**

Store at room temperature between 36-86 degrees F (2-30 degrees C) away from heat and light. Do not store in the bathroom. Keep all medicines away from children and pets.