

Palmetto State Pharmaceuticals

Patient Drug Education

CARBOFED DM SYRUP

USES:

This combination medication is used to treat symptoms caused by the common cold, flu, allergies, hay fever, or other breathing illnesses (e.g., sinusitis, bronchitis). Dextromethorphan is a cough suppressant that affects a certain part of the brain (cough center), reducing the urge to cough. Decongestants help relieve stuffy nose symptoms. Antihistamines relieve watery eyes, itchy eyes/nose/throat, runny nose, and sneezing.

HOW TO TAKE THIS MEDICATION:

HOW TO USE: If you are taking the over-the-counter product, read all directions on the product package before taking this medication. If you have any questions, consult your pharmacist. If your doctor has prescribed this medication, take it as directed. Take this medication by mouth with or without food with a full glass of water (8 ounces or 240 milliliters) or as directed by your doctor. This medication may be taken with food or milk if stomach upset occurs. If you are using the liquid form, use a medication measuring device to carefully measure the prescribed dose. Do not use a household spoon. If your liquid form is a suspension, shake the bottle well before each dose. If you are using sustained-release tablets or capsules, swallow the medication whole. Do not crush, chew, or break the tablets or capsules. Doing so can destroy the long action of the drug and may increase side effects. The dosage is based on your age, medical condition and response to therapy. If you are taking this medication for allergy or hay fever symptoms, take it regularly in order to get the most benefit from it. Improper use of this medication (abuse) may result in serious harm (e.g., brain damage, seizure, death). Do not increase your dose, take it more frequently, or use it for a longer time than directed. Inform your doctor if your condition persists or worsens.

SIDE EFFECTS:

Drowsiness, dizziness, headache, blurred vision, upset stomach, nausea, or dry mouth/nose/throat may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. To relieve dry mouth, suck on (sugarless) hard candy or ice chips, chew (sugarless) gum, drink water, or use a saliva substitute. This medication can dry up and thicken mucus in your lungs, making it more difficult to breathe and clear your lungs. To help prevent this effect, drink plenty of fluids unless otherwise directed by your doctor. If this drug is a prescription, remember that your doctor has prescribed it because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: mental/mood changes (e.g., confusion, hallucinations), ringing in the ears, shaking (tremors), trouble urinating, weakness. Tell your doctor immediately if any of these rare but very serious side effects occur: easy bruising/bleeding, chest pain, fast/slow/irregular heartbeat, seizure. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking this product, tell your doctor or pharmacist if you are allergic to it; or to tartrazine; or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (e.g., asthma, emphysema), diabetes, a certain eye problem (glaucoma), heart problems, high blood pressure, kidney problems, liver disease, seizures, stomach/intestinal problems (e.g., ulcers, blockage), overactive thyroid (hyperthyroidism), urination problems (e.g., trouble urinating due to enlarged prostate, urinary retention). This drug may make you dizzy or drowsy or cause blurred vision; use caution engaging in activities requiring alertness such as driving or using machinery. Limit the use of alcohol and certain other drowsiness-causing medications. (See also Drug Interactions.) To minimize dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. This medicine may contain aspartame. If you have phenylketonuria (PKU) or any other condition that requires you to restrict your intake of aspartame (or phenylalanine), consult your doctor or pharmacist regarding the safe use of this medicine. Liquid preparations of this product may contain sugar and/or alcohol. Caution is advised if you have diabetes, alcohol dependence, or liver disease. Ask your doctor or pharmacist about the safe use of this product. The elderly may be more sensitive to the effects of this drug, especially dizziness, drowsiness, mental/mood changes, and blood pressure changes. Do not give this medication to a child under 6 years unless directed to do so by a doctor. Children may be more sensitive to the effects of antihistamines. In young children, this medication may cause agitation/excitement instead of drowsiness. During the first 6 months of pregnancy, this medication should be used only if clearly needed. It should not be used during the last 3 months of pregnancy since antihistamines may increase the risk of serious side effects (e.g., seizures) in a newborn baby. Discuss the risks and benefits with your doctor. This medication may pass into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Do not start, stop or change the dosage of any medicine without first checking with your doctor or pharmacist. This product should not be used with the following medications because very serious (rarely fatal) interactions may occur: MAO inhibitors (e.g., furazolidone, isocarboxazid, linezolid, moclobemide, phenelzine, procarbazine, selegiline, tranylcypromine), sibutramine. If you are currently using any of these medications listed above, tell your doctor or pharmacist before starting this medication. Avoid taking MAO inhibitors within 2 weeks of starting or stopping this medication. Before using this product, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially: antispasmodics (e.g., atropine, belladonna alkaloids), beta blockers (e.g., metoprolol, atenolol), drugs for Parkinson's disease (e.g., anticholinergics such as benztropine, trihexyphenidyl), guanethidine, certain inhaled anesthetics (e.g., halothane), memantine, methyldopa, reserpine, scopolamine, tricyclic antidepressants (e.g., amitriptyline, desipramine). Tell your doctor or pharmacist if you also take drugs that cause drowsiness such as: certain antihistamines (e.g., diphenhydramine), anti-seizure drugs (e.g., carbamazepine), medicine for sleep or anxiety (e.g., alprazolam, diazepam, zolpidem), muscle relaxants, narcotic pain relievers

Palmetto State Pharmaceuticals

(e.g., codeine), psychiatric medicines (e.g., chlorpromazine, risperidone, amitriptyline, trazodone). Check the labels on all your medicines (e.g., cough-and-cold products, diet aids) because they may contain ingredients that could affect your heart rate/blood pressure or cause drowsiness. Ask your pharmacist about the safe use of those products.

NOTES:

If your doctor has prescribed this medication, do not share it with others. Keep all regular medical and laboratory appointments. Do not take this product for several days before allergy testing because test results can be affected.

MISSED DOSE:

If you are taking this product on a regular schedule and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store in a tightly closed container at room temperature between 59-86 degrees F (15-30 degrees C) away from light and moisture. Do not store in the bathroom. Do not freeze liquid forms of this medication. Keep all medicines away from children and pets.