

Palmetto State Pharmaceuticals

Patient Drug Education

CELEBREX 100MG CAPSULE

USES:

This medication is a nonsteroidal anti-inflammatory drug (NSAID), specifically a COX-2 inhibitor, which relieves pain and swelling (inflammation). It is used to treat arthritis, acute pain, and menstrual pain and discomfort. The pain and swelling relief provided by this medication helps you perform more of your normal daily activities. Celecoxib is also used to decrease growths found in the intestines (colon polyps) of persons with a family history of this condition. This drug works by blocking the enzyme in your body that makes prostaglandins. Decreasing prostaglandins helps to reduce pain and swelling.

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Read the Medication Guide provided by your pharmacist before you start using celecoxib and each time you get a refill. If you have any questions regarding the information, consult your doctor or pharmacist. Take by mouth, usually once or twice daily, or as directed by your doctor. To decrease the chance of stomach upset, this drug is best taken with food. Dosage is based on your medical condition and response to therapy. The lowest effective dosage should always be used, and only for the prescribed length of time (see also Warning section). Take this medication with a full glass of water (8 oz or 240 ml) unless your doctor directs you otherwise. Do not lie down for 30 minutes after taking this medication. In certain conditions (e.g., arthritis), it may take up to two weeks, taken regularly, before significant benefits of this drug take effect. If you are taking this drug on an "as needed" basis (not on a regular schedule), remember that pain medications work best if they are used as the first signs of pain occur. If you wait until the pain has significantly worsened, the medicine may not work as well.

SIDE EFFECTS:

Stomach upset or gas may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: severe headache, unexplained weight gain, swelling of the hands or feet, change in amount of urine, difficult/painful swallowing. This drug may rarely cause serious liver disease. If you notice any of the following highly unlikely but very serious side effects, stop taking celecoxib and consult your doctor or pharmacist immediately: yellowing eyes or skin, dark urine, persistent stomach/abdominal pain, unusual fatigue. In the unlikely event you have a serious allergic reaction to this drug, seek immediate medical attention. Symptoms of a serious allergic reaction include: rash, itching, swelling, dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking celecoxib, tell your doctor or pharmacist if you are allergic to it; or to sulfa drugs, aspirin, other NSAIDs (e.g., ibuprofen), other COX-2 inhibitors; or if you have any other allergies. This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: aspirin-sensitive asthma (a history of worsening breathing with runny/stuffy nose after taking aspirin or other NSAIDs), recent heart bypass surgery (CABG). Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney problems, liver problems, heart disease (e.g., high blood pressure, congestive heart failure), alcohol use, swelling (edema), blood disorders (anemia), serious infections, stomach problems (bleeding or ulcers), asthma, growths in the nose (nasal polyps), dehydration, poorly controlled diabetes. This medicine may cause stomach bleeding. Daily use of alcohol and tobacco, especially when combined with this medicine, may increase your risk for stomach bleeding. Limit alcohol and smoking. Consult your doctor or pharmacist for more information. Caution is advised when using this drug in the elderly because they may be more sensitive to the side effects of this medication, especially stomach bleeding and kidney effects. This medication should be used only when clearly needed during the first 6 months of pregnancy. It should not be used during the last 3 months of pregnancy. Discuss the risks and benefits with your doctor. This medication passes into breast milk and may have undesirable effects on a nursing infant. Breast-feeding is not recommended while using this drug. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription products you may use, especially of: high blood pressure drugs such as ACE inhibitors (e.g., captopril, lisinopril), oral bisphosphonates (e.g., alendronate), "blood thinners" (e.g., warfarin), corticosteroids (e.g., prednisone), fluconazole, lithium, "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene). Check all prescription and nonprescription medicine labels carefully since many contain pain relievers/fever reducers (NSAIDs such as ibuprofen, naproxen, or aspirin) which are similar to this drug and if taken together may increase your risk for side effects. Low-dose aspirin, as prescribed by your doctor for specific medical reasons such as heart attack or stroke prevention (usually these dosages are 81-325 mg per day), should be continued. Consult your doctor or pharmacist for more details. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES:

Do not share this medication with others. Laboratory and/or medical tests (e.g., complete blood count, liver and kidney function tests) may be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details. Non-drug treatment for arthritis that is approved by your doctor (e.g., weight loss if needed, strengthening and conditioning exercises) may help improve your flexibility, range of motion, and joint function. Consult your doctor for specific instructions.

Palmetto State Pharmaceuticals

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store at room temperature at 77 degrees F (25 degrees C) away from light and moisture. Brief storage between 59-86 degrees F (15-30 degrees C) is permitted. Do not store in the bathroom. Keep all medicines away from children and pets.