

# Palmetto State Pharmaceuticals

---

## *Patient Drug Education*

---

---

### **CLONAZEPAM 1 MG TABLET**

---

#### **USES:**

Clonazepam is used to treat seizure disorders and panic attacks. It belongs to a class of medications called benzodiazepines which act on the brain and nerves (central nervous system) to produce a calming effect. This drug works by enhancing the effects of a certain natural chemical in the body (GABA).

#### **HOW TO TAKE THIS MEDICATION:**

**HOW TO USE:** Take this medication by mouth 2 or 3 times a day, or as directed by your doctor. Dosage is based on your medical condition and response to therapy. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same times each day. Use this medication exactly as prescribed. Do not increase your dose, take it more frequently or use it for a longer period of time than prescribed because this drug can be habit-forming. Do not suddenly stop using this drug without your doctor's approval. Some conditions may become worse when the drug is abruptly stopped. Your dose may need to be gradually decreased. When used for an extended period, this medication may not work as well and may require different dosing. Talk with your doctor if this medication stops working well. Inform your doctor if your condition persists or worsens.

#### **SIDE EFFECTS:**

Drowsiness, dizziness, blurred vision, muscle pain, excessive hair growth or loss, change in sexual desire, weight change, increased saliva production, sore gums, or cough may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these serious side effects occur: loss of coordination, mental/mood/behavior changes. Tell your doctor immediately if any of these unlikely but serious side effects occur: slowed breathing, easy bruising/bleeding, unusual eye movements, unusual fatigue, fast/pounding heartbeat, unusual muscle movement, one-sided weakness, slurred speech, trouble urinating. If you have several different types of seizure disorders, you may experience a worsening of seizure activity at the start of using this medication. Consult your doctor immediately if this happens. The addition or dosage adjustment of other anti-seizure medications may be necessary to keep seizures under control. A serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

#### **PRECAUTIONS:**

Before taking clonazepam, tell your doctor or pharmacist if you are allergic to it; or to other benzodiazepines (e.g., diazepam, lorazepam); or if you have any other allergies. This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: narrow angle glaucoma, severe liver disease. Before using this medication, tell your doctor or pharmacist your medical history, especially of: alcohol use, drug dependency, kidney disease, liver disease, lung/breathing disorders, mental/mood disorders, open angle glaucoma. This drug may make you dizzy or drowsy; use caution engaging in activities requiring alertness such as driving or using machinery. Avoid alcoholic beverages. Caution is advised when using this drug in the elderly because they may be more sensitive to its effects, especially drowsiness and confusion. Caution is advised when using this drug in children because the long-term effects on physical and mental/behavioral development are uncertain. Discuss the risks and benefits with your doctor. This medication is not recommended for use during pregnancy due to the potential for harm to an unborn baby. If you become pregnant or think you may be pregnant, inform your doctor immediately. Consult your doctor for more details. This drug passes into breast milk and may have undesirable effects on a nursing infant. Therefore, breast-feeding while using this medication is not recommended. Consult your doctor before breast-feeding.

#### **DRUG INTERACTIONS:**

This drug should not be used with the following medication because very serious interactions may occur: sodium oxybate. If you are currently using this medication, tell your doctor or pharmacist before starting clonazepam. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: certain antidepressants (e.g., fluoxetine, fluvoxamine, nefazodone), cimetidine, kava. Tell your doctor or pharmacist if you also take drugs that cause drowsiness such as: antihistamines that cause drowsiness (e.g., diphenhydramine), anti-anxiety drugs (e.g., diazepam), other anti-seizure drugs (e.g., carbamazepine), medicine for sleep (e.g., sedatives), muscle relaxants, narcotic pain relievers (e.g., codeine), psychiatric medicines (e.g., phenothiazines such as chlorpromazine, or tricyclics such as amitriptyline), tranquilizers. Check the labels on all your medicines (e.g., cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about the safe use of those products. Do not start or stop any medicine without doctor or pharmacist approval.

#### **NOTES:**

Do not share this medication with others. It is against the law. Laboratory and/or medical tests (e.g., liver function tests, complete blood count) should be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details.

#### **MISSED DOSE:**

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule.

# Palmetto State Pharmaceuticals

Do not double the dose to catch up.

## **STORAGE:**

Store at room temperature (77 degrees F or 25 degrees C) away from light and moisture. Brief storage between 59-86 degrees F (15-30 degrees C) is permitted. Do not store in the bathroom. Keep all medicines away from children and pets.