

Drug Info

CLOTRIMAZOLE+BETAMETHAS (LOTRISONE)

BEFORE USING THIS MEDICINE:

- * Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- * Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.

COMMON USES FOR THIS MEDICINE:

- 1- For the treatment of RINGWORM OF THE BODY (Tinea corporis) due to Epidermophyton rubrum.
- 2- For the treatment of RINGWORM OF THE FOOT (Tinea pedis; Athlete's foot) due to Epidermophyton rubrum.
- 3- For the treatment of RINGWORM OF THE GROIN (Tinea cruris; Jock itch) due to Epidermophyton rubrum.
- 4- For the treatment of MYCOSES, UNSPECIFIED (SYSTEMIC, SKIN & EYE FUNGAL INFECTION)
- 5- For the treatment of PRURITUS AND RELATED CONDITIONS

HOW TO USE THIS MEDICINE:

Use this medicine exactly as directed on the label, unless instructed differently by your doctor

- * CLEANSE & DRY area first before applying.
- * Use this medicine for the full prescribed treatment period.
- * Keep this medicine away from the EYES, NOSE, MOUTH, & other mucous membranes. Do not use OCCLUSIVE dressing unless directed to do so by your doctor.
- * This medication is for EXTERNAL USE only.

WHAT TO DO IF YOU MISS A DOSE:

Apply as soon as possible when remembered unless time for next dose. No double doses.

HOW TO STORE THIS MEDICINE:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

STOP TAKING THIS MEDICINE AND GET EMERGENCY HELP IMMEDIATELY IF ANY OF THE FOLLOWING OCCURS (SEE SYMPTOMS BELOW):

Shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching.

CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Skin rash; Skin irritation; Weight gain; Muscle weakness; Menstrual irregularities; Swelling of feet or lower legs; Swelling usually of face or lips; Severe skin itching with patches.

OTHER SIDE EFFECTS:

Headache; Nausea; Gastric or intestinal ulcers; Difficulty falling asleep; Vertigo; High blood pressure; Unusual tiredness or weakness; Abnormal hair growth; Low blood pressure; Increased eye pressure; Electrolyte imbalance; Glaucoma; Vomiting; Increased appetite.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

PRECAUTIONS AND WARNINGS:

- * Safety of use during pregnancy has not been established. This medicine SHOULD NOT BE USED DURING PREGNANCY unless benefit justifies potential risk to the fetus. CALL YOUR DOCTOR.
- * Notify your doctor if condition worsens or shows no improvement.