

# Drug Info

## CODEINE + APAP (TYLENOL W/COD.)

### BEFORE USING THIS MEDICINE:

- \* Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- \* Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.
- \* Inform your doctor if you have any of these conditions : KIDNEY DISEASE; CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD); ASTHMA; LIVER PROBLEMS.

### COMMON USES FOR THIS MEDICINE:

- 1- For the treatment of PAIN, GENERALIZED

### HOW TO USE THIS MEDICINE:

Use this medicine exactly as directed on the label, unless instructed differently by your doctor

- \* This medicine can be taken with or without food, REGARDLESS of meal times. If stomach upset occurs, take with food.
- \* DO NOT EXCEED recommended dosage; excessive use may lead to adverse effects or loss of effectiveness. Do not adjust the dose unless instructed.

### WHAT TO DO IF YOU MISS A DOSE:

This medicine is usually used as needed. If you miss a dose administer when or if needed. No double doses.

### HOW TO STORE THIS MEDICINE:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

### CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Yellowing of eyes or skin; Jerkings of face, trunk, or extremities; Shortness of breath; Skin rash; Irregular heartbeat; Low blood pressure; Hot sensation and redness of face.

### OTHER SIDE EFFECTS:

Sedation; Nausea; Constipation; Dizziness; Vomiting; Dry mouth; Sweating; Anemia; Abnormal blood cell count; Changes in urination frequency or amount; Decreased appetite; Blurred vision; Confusion; Headache; Lightheadedness; Feeling of general discomfort; Drowsiness.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

### PRECAUTIONS AND WARNINGS:

- \* Safety of use during pregnancy has not been established. This medicine SHOULD NOT BE USED DURING PREGNANCY unless benefit justifies potential risk to the fetus. CALL YOUR DOCTOR.
- \* This medicine may cause drowsiness, dizziness, or blurred vision. DO NOT drive or operate machines until you know how you react to this medicine.
- \* AVOID concurrent use of alcohol or other CNS depressants as it may intensify the drowsiness, dizziness, or lightheadedness effect of this medicine.
- \* This medicine may cause dizziness, lightheadedness or fainting. Avoid sudden changes in posture. BE CAREFUL and slowly sit up or stand up.
- \* Prolonged use of this medication may cause physical and psychological dependence, do not exceed recommended dose.
- \* Strictly limit alcohol consumption. LIVER PROBLEMS may be more likely to occur if you drink alcoholic beverages regularly while taking this medicine.