

Drug Info

CODEINE + GUAIFENESIN (ROBITUSSIN-AC)

BEFORE USING THIS MEDICINE:

- * Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- * Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.
- * Inform your doctor if you have any of these conditions : KIDNEY DISEASE; LIVER PROBLEMS; URINARY RETENTION.

COMMON USE FOR THIS MEDICINE:

1- For the relief of COUGH

HOW TO USE THIS MEDICINE:

- Use this medicine exactly as directed on the label, unless instructed differently by your doctor
- * Take this medicine WITH food, a meal, or a snack. Do not take on an empty stomach.

WHAT TO DO IF YOU MISS A DOSE:

Take when remembered unless time for next dose. No double doses.

HOW TO STORE THIS MEDICINE:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Skin rash; Shortness of breath; Irregular heartbeat; Low blood pressure; Hot sensation and redness of face.

OTHER SIDE EFFECTS:

Dizziness; Vomiting; Nausea; Headache; Stomach discomfort or pain; Constipation; Sedation; Dry mouth; Sweating.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

PRECAUTIONS AND WARNINGS:

- * Safety of use during pregnancy has not been established. This medicine SHOULD NOT BE USED DURING PREGNANCY unless benefit justifies potential risk to the fetus. CALL YOUR DOCTOR.
- * This medicine may cause drowsiness, dizziness, or blurred vision. DO NOT drive or operate machines until you know how you react to this medicine.
- * This medicine may cause dizziness, lightheadedness or fainting. Avoid sudden changes in posture. BE CAREFUL and slowly sit up or stand up.
- * AVOID concurrent use of alcohol or other CNS depressants as it may intensify the drowsiness, dizziness, or lightheadedness effect of this medicine.
- * Because of INTERACTIONS, report the use of any other prescription or nonprescription medicines, including natural/herbal remedies, to your doctor.
- * The results of some MEDICAL or LAB tests may be affected by this medicine. Inform the doctor in charge that you are taking this medicine.