

Palmetto State Pharmaceuticals

Patient Drug Education

DIAZEPAM 10 MG TABLET

USES:

Diazepam is used to treat anxiety, acute alcohol withdrawal, and seizures. It is also used to relieve muscle spasms and to provide sedation before medical procedures. This medication belongs to a class of drugs called benzodiazepines which act on the brain and nerves (central nervous system) to produce a calming effect. It works by enhancing the effects of a certain natural chemical in the body (GABA).

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Take this medication by mouth as directed by your doctor. The dosage is based on your medical condition and response to therapy. If you are using the oral concentrate solution, use the dropper provided to carefully measure out the dose and mix it in liquid or soft food (e.g., applesauce, pudding) just before taking. Use this medication exactly as prescribed. Do not increase your dose, take it more frequently or use it for a longer period of time than prescribed because this drug can be habit-forming. Also, if used for an extended period of time or for seizure control, do not suddenly stop using this drug without your doctor's approval. Some conditions may become worse when the drug is abruptly stopped. Your dose may need to be gradually decreased. When used for an extended period, this medication may not work as well and may require different dosing. Talk with your doctor if this medication stops working well. Avoid eating grapefruit or drinking grapefruit juice while being treated with this medication unless your doctor instructs you otherwise. Grapefruit juice can increase the amount of certain medications in your bloodstream. Consult your doctor or pharmacist for more details. Inform your doctor if your condition persists or worsens.

SIDE EFFECTS:

Drowsiness, dizziness, fatigue, constipation, blurred vision, or headache may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: mental/mood changes, slurred speech, clumsiness, trouble walking, decreased/increased interest in sex, tremor, trouble urinating, sleep disturbances. Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: stomach/abdominal pain, persistent nausea, vomiting, yellowing eyes or skin, dark urine, persistent sore throat or fever. A serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking diazepam, tell your doctor or pharmacist if you are allergic to it; or to other benzodiazepines (e.g., alprazolam, lorazepam); or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease, lung/breathing problems (e.g., COPD, sleep apnea), drug or alcohol abuse. This drug may make you dizzy, drowsy or cause blurred vision; use caution engaging in activities requiring alertness such as driving or using machinery. Limit alcoholic beverages. Caution is advised when using this drug in the elderly because they may be more sensitive to the effects of the drug, especially the drowsiness effect. This medication is not recommended for use during pregnancy due to the potential for harm to an unborn baby. If you become pregnant or think you may be pregnant, inform your doctor immediately. Consult your doctor for more details. This drug passes into breast milk and may have undesirable effects on a nursing infant. Therefore, breast-feeding while using this medication is not recommended. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

See also the How to Use section. This drug should not be used with the following medications because very serious interactions may occur: fluvoxamine, sodium oxybate. If you are currently using any of these medications, tell your doctor or pharmacist before starting diazepam. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: antacids, certain anti-depressants (e.g., fluoxetine, nefazodone), cimetidine, clozapine, digoxin, disulfiram, kava, levodopa. Tell your doctor or pharmacist if you also take drugs that cause drowsiness such as: antihistamines that cause drowsiness (e.g., diphenhydramine), anti-seizure drugs (e.g., carbamazepine), medicine for sleep (e.g., sedatives), muscle relaxants, narcotic pain relievers (e.g., codeine), psychiatric medicines (e.g., phenothiazines such as chlorpromazine, or tricyclics such as amitriptyline), tranquilizers. Check the labels on all your medicines (e.g., cough-and-cold products) because they may contain drowsiness-causing ingredients. Ask your pharmacist about the safe use of those products. This product can affect the results of certain lab tests. Make sure laboratory personnel and your doctors know you use this drug. Smoking can decrease the effectiveness of this drug (through liver enzyme induction). Tell your doctor if you smoke or if you have recently stopped smoking because your dose may need to be adjusted. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES:

Do not share this medication with others. It is against the law. If this drug is used for an extended period of time, laboratory and/or medical tests (e.g., liver function tests, complete blood count) may be performed periodically to check for side effects. Consult your doctor for more details.

Palmetto State Pharmaceuticals

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up. If you are taking this medication for seizures, take it if remembered within 1 hour of the missed dose, but skip it if more than 1 hour has passed.

STORAGE:

Store at room temperature between 59 and 86 degrees F (15-30 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets.