

Palmetto State Pharmaceuticals

Patient Drug Education

ETODOLAC 400 MG TABLET SA

USES:

This medication is used to reduce pain, swelling, and joint stiffness from arthritis. Etodolac is known as a nonsteroidal anti-inflammatory drug (NSAID).

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Read the Medication Guide provided by your pharmacist before you start using etodolac and each time you get a refill. If you have any questions regarding the information, consult your doctor or pharmacist. Take this medication by mouth usually once daily with a full glass of water (8 ounces or 240 milliliters), unless your doctor directs you otherwise. Do not lie down for at least 30 minutes after taking this drug. Swallow this medication whole. Do not crush, chew, or break the tablets; doing so can destroy the long action of the drug and may increase side effects. If stomach upset occurs while taking this medication, take it with food, milk, or an antacid. The dosage is based on your medical condition and response to therapy. Do not increase your dose or take it more frequently than recommended because this may increase your risk of stomach bleeding. It may take up to two weeks, taken regularly, before the full benefits of this drug take effect. Inform your doctor if your condition worsens.

SIDE EFFECTS:

Upset stomach, diarrhea, gas, nausea, vomiting, constipation, fatigue, or dizziness may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: stomach pain, swelling of the hands or feet, sudden or unexplained weight gain, vision changes, hearing changes (e.g., ringing in the ears), mental/mood changes, fast/pounding heartbeat, persistent/severe headache, fainting, difficult/painful swallowing. Tell your doctor immediately if any of these rare but very serious side effects occur: change in the amount of urine, easy bruising/bleeding, signs of infection (e.g., fever, persistent sore throat), unexplained stiff neck. This drug may rarely cause serious (possibly fatal) liver disease. If you notice any of the following highly unlikely but very serious side effects, stop taking etodolac and consult your doctor or pharmacist immediately: yellowing eyes or skin, dark urine, unusual/extreme tiredness, severe stomach/abdominal pain, persistent nausea/vomiting. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking etodolac, tell your doctor or pharmacist if you are allergic to it; or to aspirin or other NSAIDs (e.g., ibuprofen, naproxen, celecoxib); or if you have any other allergies. This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: aspirin-sensitive asthma (a history of worsening breathing with runny/stuffy nose after taking aspirin or other NSAIDs), recent heart bypass surgery (CABG). Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, poorly controlled diabetes, stomach/intestinal problems (e.g., bleeding, ulcers), heart disease (e.g., congestive heart failure, history of heart attack), stroke, high blood pressure, swelling (edema, fluid retention), dehydration, blood disorders (e.g., anemia), bleeding or clotting problems, asthma, growths in the nose (nasal polyps). Before having surgery, tell your doctor or dentist that you are taking this medication. This drug may make you dizzy; use caution engaging in activities requiring alertness such as driving or using machinery. This medicine may cause stomach bleeding. Daily use of alcohol and tobacco, especially when combined with this medicine, may increase your risk for stomach bleeding. Limit alcohol and stop smoking. Consult your doctor or pharmacist for more information. This medication may make you more sensitive to the sun. Avoid prolonged sun exposure, tanning booths or sunlamps. Use a sunscreen and wear protective clothing when outdoors. Caution is advised when using this drug in the elderly because they may be more sensitive to its side effects, especially stomach bleeding and kidney effects. Etodolac should be used only when clearly needed during the first 6 months of pregnancy. It is not recommended for use during the last 3 months of pregnancy due to possible harm to an unborn baby and interference with normal labor/delivery. Discuss the risks and benefits with your doctor. It is not known if this medication passes into breast milk. Breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Your healthcare professionals (e.g., doctor or pharmacist) may already be aware of any possible drug interactions and may be monitoring you for it. Do not start, stop, or change the dosage of any medicine before checking with them first. This drug should not be used with the following medications because very serious interactions may occur: cidofovir, ketorolac. If you are currently using any of these medications listed above, tell your doctor or pharmacist before starting etodolac. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: anti-platelet drugs (e.g., cilostazol, clopidogrel), oral bisphosphonates (e.g., alendronate), "blood thinners" (e.g., enoxaparin, heparin, warfarin), corticosteroids (e.g., prednisone), cyclosporine, digoxin, high blood pressure drugs (including ACE inhibitors such as captopril, angiotensin II receptor antagonists such as losartan, and beta-blockers such as metoprolol), lithium, methotrexate, pemetrexed, probenecid, SSRI antidepressants (e.g., fluoxetine, sertraline), "water pills" (diuretics such as furosemide, hydrochlorothiazide, triamterene). Check all prescription and nonprescription medicine labels carefully since many contain pain relievers/fever reducers (NSAIDs such as aspirin, ibuprofen or naproxen) which are similar to this drug and if taken together may increase your risk for side effects. Low-dose aspirin, as prescribed by your doctor for specific medical reasons such as heart attack or stroke prevention (usually these dosages are 81-325 milligrams per day), should be continued. Consult your doctor or pharmacist for more details. This product can affect the results of certain lab tests (urine tests). Make sure laboratory personnel and your doctors know

Palmetto State Pharmaceuticals

you use this drug.

NOTES:

Do not share this medication with others. Laboratory and/or medical tests (e.g., complete blood count, liver and kidney function tests) may be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details. Non-drug treatment for arthritis as approved by your doctor (e.g., weight loss if needed, strengthening and conditioning exercises) may help improve your flexibility (range of motion) and improve your joint function. Consult your doctor for specific instructions.

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store at room temperature between 68-77 degrees F (20-25 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets.