

Palmetto State Pharmaceuticals

Patient Drug Education

GABAPENTIN 800 MG TABLET

USES:

Gabapentin is used with other medications to help control seizures in adults and children (3 years of age and older). It is also used to relieve nerve pain associated with shingles (herpes zoster) infection in adults.

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Take this medication by mouth, with or without food, as directed by your doctor. The dosage is based on your medical condition and response to therapy. During the first few days, your doctor may gradually increase your dose so your body can adjust to the medication. To minimize side effects, take the very first dose at bedtime. Use this medication regularly in order to get the most benefit from it. This drug works best when the amount of medicine in your body is kept at a constant level. Therefore it is best to take gabapentin at evenly spaced intervals throughout the day and night. Do not take this medication more often or increase your dose without consulting your doctor. Your condition will not improve any faster and the risk of serious side effects may be increased. Do not stop taking this medication without consulting your doctor. Some conditions such as seizures may become worse when the drug is abruptly stopped. Your dose may need to be gradually decreased. Antacids containing aluminum or magnesium may interfere with the absorption of this medication. Therefore it is best to take gabapentin at least 2 hours after taking an antacid. Inform your doctor if your condition does not improve or worsens.

SIDE EFFECTS:

Drowsiness, dizziness, unsteadiness, fatigue, vision changes, weight gain, nausea, dry mouth, or constipation may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these serious side effects occur: tremor or shaking, swollen arms/legs. Tell your doctor immediately if any of these unlikely but serious side effects occur: loss of coordination, mental/mood changes. Tell your doctor immediately if any of these highly unlikely but very serious side effects occur: persistent sore throat/fever/cough, unusual bleeding/bruising, pain/redness of arms or legs, chest pain, trouble breathing, fast/slow/irregular heartbeat, hearing loss, stomach/abdominal pain. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking gabapentin, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease. This drug may make you dizzy or drowsy; use caution engaging in activities requiring alertness such as driving or using machinery. Limit alcoholic beverages as they may worsen these effects. Caution is advised when using this drug in the elderly because they may be more sensitive to its effects, especially swollen arms/legs or loss of coordination. Caution is advised when using this drug in children because they may be more sensitive to its effects, especially the mental/mood changes (e.g., hostility). This medication should be used only when clearly needed during pregnancy. Discuss the risks and benefits with your doctor. Gabapentin passes into breast milk. Because the effects of this drug on the nursing infant are unknown, consult your doctor before breast-feeding.

DRUG INTERACTIONS:

See also How to Use section. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: antacids containing aluminum or magnesium, morphine. This product can affect the results of certain lab tests for urine protein. Make sure laboratory personnel and your doctors know you use this drug. Do not start or stop any medicine without doctor or pharmacist approval.

NOTES:

Do not share this medication with others.

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store at room temperature (77 degrees F or 25 degrees C) away from light and moisture. Brief storage between 59-86 degrees F (15-30 degrees C) is permitted. Do not store in the bathroom. Keep all medicines away from children and pets.