

Palmetto State Pharmaceuticals

Patient Drug Education

INDOMETHACIN 25 MG CAPSULE

USES:

Indomethacin is used to relieve pain, swelling, and joint stiffness caused by arthritis, gout, bursitis, and tendonitis. Reducing these symptoms helps you do more of your normal daily activities. This medication is known as a nonsteroidal anti-inflammatory drug (NSAID).

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Read the Medication Guide provided by your pharmacist before you start using indomethacin and each time you get a refill. If you have any questions regarding the information, consult your doctor or pharmacist. Take this medication by mouth with a full glass of water (8 ounces or 240 milliliters) unless your doctor directs you otherwise. Do not lie down for at least 30 minutes after taking this drug. Take this medication either with food, right after meals, or with antacids to prevent stomach upset. Dosage is based on your medical condition and response to therapy. Do not take more than 200 milligrams per day. To minimize side effect risks (e.g., stomach bleeding), use this medication at the lowest effective dose for the shortest possible length of time. Do not increase your dose or take it more often than prescribed. For ongoing conditions such as arthritis, continue taking it as directed by your doctor. Discuss the risks and benefits with your doctor or pharmacist. In certain conditions (e.g., arthritis), it may take up to 4 weeks of regular use before the full benefits of this drug take effect. If you are taking this drug on an "as needed" basis (not on a regular schedule), remember that pain medications work best if they are used as the first signs of pain occur. If you wait until the pain has significantly worsened, the medicine may not work as well. Inform your doctor if your condition worsens.

SIDE EFFECTS:

See also Warning section. Upset stomach, heartburn, headache, tiredness, drowsiness, or dizziness may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: stomach pain, swelling of the hands/feet, sudden/unexplained weight gain, vision changes, hearing changes (e.g., ringing in the ears), mental/mood changes (e.g., confusion, hallucinations), fast/pounding heartbeat, persistent/severe headache, fainting, muscle weakness, uncontrollable movements, difficult/painful swallowing. Tell your doctor immediately if any of these rare but very serious side effects occur: change in the amount of urine, easy bruising/bleeding, signs of infection (e.g., fever, persistent sore throat), unexplained stiff neck, seizures. This drug may rarely cause serious (possibly fatal) liver disease. If you notice any of the following rare but very serious side effects, stop taking indomethacin and tell your doctor immediately: yellowing eyes/skin, dark urine, unusual/extreme tiredness, severe stomach/abdominal pain, persistent nausea/vomiting. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking indomethacin, tell your doctor or pharmacist if you are allergic to it; or to aspirin or other NSAIDs (e.g., ibuprofen, naproxen, celecoxib); or if you have any other allergies. This medication should not be used if you have certain medical conditions. Before using this medicine, consult your doctor or pharmacist if you have: aspirin-sensitive asthma (a history of worsening breathing with runny/stuffy nose after taking aspirin or other NSAIDs), severe kidney disease, recent heart bypass surgery (CABG). Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, poorly controlled diabetes, stomach/intestine/esophagus problems (e.g., bleeding, ulcers), heart disease (e.g., congestive heart failure, history of heart attack), blood disorders (e.g., anemia), bleeding or clotting problems, high blood pressure, swelling (edema, fluid retention), a severe loss of body water (dehydration), stroke, mental/mood conditions (e.g., depression, psychoses), seizures, Parkinson's disease, high level of potassium in the blood, asthma, growths in the nose (nasal polyps). Before having surgery, tell your doctor or dentist that you are using this medication. This drug may make you dizzy or drowsy; use caution engaging in activities requiring alertness such as driving or using machinery. This medicine may cause stomach bleeding. Daily use of alcohol and tobacco may increase your risk for stomach bleeding, especially when combined with this medicine. Limit alcohol and stop smoking. Consult your doctor or pharmacist for more information. This medication may make you more sensitive to the sun. Avoid prolonged sun exposure, tanning booths, and sunlamps. Use a sunscreen and wear protective clothing when outdoors. The elderly may be more sensitive to the side effects of this drug, especially stomach/intestinal bleeding, kidney effects, and mental/mood changes. Children may be more sensitive to the side effects of this drug, especially serious liver problems. Caution is advised when this drug is used in children. Discuss the risks and benefits of treatment with your doctor. This medication should be used only when clearly needed during the first 6 months of pregnancy. It is not recommended for use during the last 3 months of pregnancy due to possible harm to the unborn baby and interference with normal labor/delivery. Discuss the risks and benefits with your doctor. This drug passes into breast milk. Therefore, breast-feeding while using this drug is not recommended. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Your healthcare professionals (e.g., doctor or pharmacist) may already be aware of any possible drug interactions and may be monitoring you for it. Do not start, stop or change the dosage of any medicine before checking with them first. This drug should not be used with the following medications because very serious interactions may occur: cidofovir, other NSAIDs (e.g., diflunisal, ketorolac), high doses of aspirin and related drugs (salicylates), a certain "water pill" (a diuretic called triamterene). If you are currently using any of these medications listed above, tell your doctor or pharmacist before starting indomethacin. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use,

Palmetto State Pharmaceuticals

especially of: anti-platelet drugs (e.g., cilostazol, clopidogrel), oral bisphosphonates (e.g., alendronate), "blood thinners" (e.g., enoxaparin, heparin, warfarin), corticosteroids (e.g., prednisone), cyclosporine, digoxin, high blood pressure drugs (including ACE inhibitors such as captopril, angiotensin receptor blockers such as losartan, and beta-blockers such as metoprolol), lithium, methotrexate, pemetrexed, potassium supplements, probenecid, SSRI antidepressants (e.g., fluoxetine, sertraline), other "water pills" (diuretics such as furosemide, hydrochlorothiazide, spironolactone). Check all prescription and nonprescription medicine labels carefully since many contain pain relievers/fever reducers (NSAIDs such as aspirin, ibuprofen or naproxen) which are similar to this drug and if taken together may increase your risk for side effects. Low-dose aspirin, as prescribed by your doctor for specific medical reasons such as heart attack or stroke prevention (usually at dosages of 81-325 milligrams per day), should be continued. Consult your doctor or pharmacist for more details. This medication can affect the results of certain lab tests. Make sure laboratory personnel and your doctors know you use this drug.

NOTES:

Do not share this medication with others. Laboratory and/or medical tests (e.g., complete blood count, liver and kidney function tests) may be performed periodically to monitor your progress or check for side effects. Consult your doctor for more details. Non-drug treatment for arthritis that is approved by your doctor (e.g., weight loss if needed, strengthening and conditioning exercises) may help improve your flexibility, range of motion, and joint function. Consult your doctor for specific instructions.

MISSED DOSE:

If you are prescribed this drug on a regular schedule (not just "as needed") and you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

STORAGE:

Store at room temperature between 59-86 degrees F (15-30 degrees C) away from light and moisture. Do not store in the bathroom. Keep all medicines away from children and pets.