

Palmetto State Pharmaceuticals

Patient Drug Education

LYRICA 100 MG CAPSULE

USES:

This medication is used to treat pain caused by nerve damage due to diabetes and shingles (herpes zoster) infection. It is also used with other medications to treat certain types of seizures (partial onset seizures).

HOW TO TAKE THIS MEDICATION:

HOW TO USE: Read the Patient Information Leaflet provided by your pharmacist before you start using pregabalin and each time you get a refill. If you have any questions regarding the information, consult your doctor or pharmacist. Take this medication by mouth, usually twice or three times a day, or as directed by your doctor. You may take it with or without food. When you start this medication, your dosage will probably need to be increased slowly by your doctor to reduce side effects, especially dizziness or drowsiness. Your dosage is based on your medical condition and response to therapy. Use this medication regularly in order to get the most benefit from it. This drug works best when the amount of medicine in your body is kept at a constant level. Therefore it is best to take pregabalin at evenly spaced intervals throughout the day and night. Do not stop taking this drug suddenly without your doctor's approval since seizures may come back, or you may have a withdrawal reaction. This medication may cause dependence, especially if it has been used regularly for an extended period of time, or if it has been used in high doses. In such cases, if you suddenly stop this drug, withdrawal reactions may occur. Such reactions can include difficulty sleeping, nausea, headache and diarrhea. Report to your doctor immediately any such reactions. When stopping extended, regular treatment with this drug, gradually reducing the dosage as directed will help prevent withdrawal reactions. Consult your doctor or pharmacist for more details. Though it is very unlikely to occur, this medication can also result in abnormal drug-seeking behavior (addiction/habit-forming). Do not increase your dose, take it more frequently or use it for a longer period of time than prescribed. Properly stop the medication when so directed. This will lessen the chances of becoming addicted. Inform your doctor if your condition persists or worsens.

SIDE EFFECTS:

Drowsiness, dizziness, confusion, difficulty concentrating, unsteadiness, fatigue, swollen arms/legs, increased appetite, weight gain, nausea, dry mouth or constipation may occur. If any of these effects persist or worsen, notify your doctor or pharmacist promptly. Remember that your doctor has prescribed this medication because the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor immediately if any of these unlikely but serious side effects occur: change in amount of urine, difficulty speaking, loss of coordination, mental/mood changes, muscle pain/tenderness/weakness (especially if you are tired or have a fever), stomach/abdominal pain, uncontrolled movements (e.g., tremor, twitching), unusual tiredness, vision changes. Tell your doctor immediately if any of these rare but very serious side effects occur: unusual bleeding/bruising, slow or irregular heartbeat. A very serious allergic reaction to this drug is unlikely, but seek immediate medical attention if it occurs. Symptoms of a serious allergic reaction may include: rash, itching, swelling, severe dizziness, trouble breathing. If you notice other effects not listed above, contact your doctor or pharmacist.

PRECAUTIONS:

Before taking pregabalin, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. Before using this medication, tell your doctor or pharmacist your medical history, especially of: diabetes, heart problems (e.g., heart failure), kidney disease. This drug may make you dizzy or drowsy or cause blurred vision; use caution engaging in activities requiring alertness such as driving or using machinery. Avoid alcoholic beverages because they may increase the risk of this drug's side effects. Kidney function declines as you grow older. This medication is removed by the kidneys. Caution is advised when using this drug in the elderly because they may be more sensitive to its effects, especially drowsiness. This medication should be used only when clearly needed during pregnancy. Discuss the risks and benefits with your doctor. It is not known whether this drug passes into breast milk. Consult your doctor before breast-feeding.

DRUG INTERACTIONS:

Your healthcare professionals (e.g., doctor or pharmacist) may already be aware of any possible drug interactions and may be monitoring you for it. Do not start, stop or change the dosage of any medicine before checking with them first. Before using this medication, tell your doctor or pharmacist of all prescription and nonprescription/herbal products you may use, especially of: certain diabetic medications (glitazones such as rosiglitazone). Tell your doctor or pharmacist if you also take drugs that cause drowsiness such as: certain antihistamines (e.g., diphenhydramine), anti-anxiety drugs (e.g., diazepam), anti-seizure drugs (e.g., carbamazepine), medicine for sleep (e.g., sedatives), muscle relaxants, narcotic pain relievers (e.g., codeine), psychiatric medicines (e.g., phenothiazines such as chlorpromazine, or tricyclics such as amitriptyline), tranquilizers. Check the labels on all your medicines (e.g., cough-and-cold products) because they may contain drowsiness-causing ingredients. Ask your pharmacist about the safe use of those products.

NOTES:

Do not share this medication with others. It is against the law. Laboratory tests may be done to measure kidney function.

MISSED DOSE:

If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule.

Palmetto State Pharmaceuticals

Do not double the dose to catch up.

STORAGE:

Store at room temperature at 77 degrees F (25 degrees C) away from light and moisture. Brief storage between 59 - 86 degrees F (15-30 degrees C) is permitted. Do not store in the bathroom. Keep all medicines away from children and pets.