

Drug Info

NEOMYCIN-POLYMYXIN-HC TOP. (CORTISPORIN)

BEFORE USING THIS MEDICINE:

- * Inform your Doctor or Pharmacist if you are allergic to any prescription or non-prescription medicine; taking other medicines or have other medical problems; pregnant, planning a pregnancy, or breastfeeding.
- * Check with your Doctor, Pharmacists, or Nurse if information in this leaflet causes you special concern or if you want additional information about your medicine and its use.

COMMON USE FOR THIS MEDICINE:

- 1- For the treatment of LOCAL INFECTIONS OF SKIN AND SUBCUTANEOUS TISSUE
- 2- For the treatment of PRURITUS AND RELATED CONDITIONS

HOW TO USE THIS MEDICINE:

Use this medicine exactly as directed on the label, unless instructed differently by your doctor

- * WASH hands prior to administration.
- * CLEANSE & DRY area first before applying.
- * WASH hands immediately after using medication.
- * Keep this medicine away from the EYES, NOSE, MOUTH, & other mucous membranes.
- * This medication is for EXTERNAL USE only.

WHAT TO DO IF YOU MISS A DOSE:

Apply as soon as possible when remembered unless time for next dose. No double doses.

HOW TO STORE THIS MEDICINE:

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

STOP TAKING THIS MEDICINE AND GET EMERGENCY HELP IMMEDIATELY IF ANY OF THE FOLLOWING OCCURS (SEE SYMPTOMS BELOW):

Shortness of breath, fast or irregular breathing, fever, joint pain, fast pulse, lightheadedness or fainting, swelling of face, hives, & itching.

CALL YOUR DOCTOR IMMEDIATELY IF YOU EXPERIENCE:

Weight gain; Muscle weakness; Hearing loss, ear ringing, unsteadiness; Menstrual irregularities; Changes in urination frequency or amount; Swelling of feet or lower legs; Dizziness; Skin rash; Jerkings of face, trunk, or extremities; Decreased appetite; Drowsiness; Difficulty breathing; Itching; Nausea; Swelling; Increased thirst; Vomiting; Swelling usually of face or lips; Severe skin itching with patches.

OTHER SIDE EFFECTS:

Diarrhea; Headache; Gastric or intestinal ulcers; Difficulty falling asleep; Vertigo; High blood pressure; Unusual tiredness or weakness; Abnormal hair growth; Low blood pressure; Increased eye pressure; Electrolyte imbalance; Glaucoma; Increased appetite.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

PRECAUTIONS AND WARNINGS:

- * This medicine is CONTRAINDICATED for use during pregnancy. CALL YOUR DOCTOR IMMEDIATELY IF PREGNANT or intend to become pregnant.
- * DO NOT place bandages, dressing, cosmetics, or other skin products over the affected area unless directed by your doctor.
- * Notify your doctor if condition worsens or shows no improvement.